

ACCEPTED  
VIA NOTIFICATION  
LABEL NOT REVIEWED

DOC ID: 555909

March 9, 2018

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration



**Onslaught**

## Microencapsulated Insecticide

**ACTIVE INGREDIENT:**

(S)-cyano(3-phenoxyphenyl)methyl-(S)-4-chloro-alpha-(1-methylethyl)benzeneacetate .....	6.40%
*OTHER INGREDIENTS .....	93.60%
	100.00%

\*Contains petroleum distillates

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**FIRST AID**

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**NOTE TO PHYSICIAN:** Contains petroleum distillate - vomiting may cause aspiration pneumonia. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

Manufactured for: **MGK**, 8810 Tenth Avenue North, Minneapolis, MN 55427  
EPA Reg. No. 1021-1815  
F28371-107  
Rev. 0616-1017

EPA Est. No. 1021-MN-2<sup>A</sup>, 70815-GA-1<sup>C</sup>, 58996-MO-1<sup>D</sup>, 39578-TX-1<sup>F</sup>,  
228-IL-002<sup>S</sup>, 10508-OH-001<sup>K</sup>, 67572-GA-1<sup>L</sup>, 63416-WI-001<sup>M</sup>, 5905-GA-01<sup>Q</sup>  
Superscript is the first character of the lot number.

**NET CONTENTS: 1 GALLON**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear protective eyewear (goggles, face shield or safety glasses with side shields). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and other aquatic invertebrates. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage system. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirement of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA. This product is highly toxic to bees and other pollinating insects exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees or other pollinating insects are foraging in the treatment area.


**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**See inside booklet for Directions for Use and Storage and Disposal**



PULL HERE TO OPEN / PRESS TO RESEAL

PF-54046-7

FIRST AID	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not give <b>any</b> liquid to the person.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or a doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>NOTE TO PHYSICIAN:</b> Contains petroleum distillate – vomiting may cause aspiration pneumonia.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear protective eyewear (goggles, face shield or safety glasses with side shields). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and other aquatic invertebrates. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage system. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirement of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

This product is highly toxic to bees and other pollinating insects exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees or other pollinating insects are foraging in the treatment area.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS:**

- Remove pets and birds and cover fish aquariums before spraying.
- Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm.
- Do not apply to classrooms while in use. Do not apply in institutions (including libraries, schools, sports facilities, etc.) in the immediate area where occupants are present.
- DO NOT use this product as a fogger or apply it as a space spray.
- Do not apply in electrical conduits, junction and switch boxes, motor housings, or other electrical equipment due to shock hazard from water-based spray.

- Use only
- Remove p
- Keep all p
- DO NOT I
- DO NOT z
- water if th
- pesticides
- Do not ap
- Do not ap

**For control**  
Buildings, Th  
and Grocery  
Lawns, Trees

**Insects Con**  
Ants (excludir  
Aphids  
Asian Lady Be  
Bed Bugs  
Beetles  
Boxelder Bug  
Brown Marmc  
Carpet Beetle  
Centipedes

**Flying Insec**  
Blow Flies  
Chiggers  
Clothes Moth  
Cluster Flies

**Biting and S**  
Bed Bugs  
Bees  
Biting Flies  
Chiggers  
Deer Flies

**Stored Prod**  
Angoumois Gr  
Ants  
Caddises  
Cigarette Beetl  
Confused Flou  
Dark Mealworm  
Driedfruit Beet  
Drugstore Bee

emergencies or  
 field or safety  
 contaminated  
 as, gutters or  
 sticide off the  
 (this product  
 S) permit and  
 flying sewage  
 llow it to drift  
 are occupants  
 y.

- Use only in areas described.
  - Remove pets and cover fish aquariums and pet's food and water dishes before spraying.
  - Keep all people (especially children) and pets out of areas being treated, and restrict access to these areas until all surfaces are dry.
  - DO NOT leave product where children or pets can come into contact with it.
  - DO NOT allow spray to contact food or food-containing surfaces, feed, or water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated with this product. Food utensils such as teaspoons and measuring cups must not be used for food purposes after use with pesticides.
  - Do not apply when food processing facility is in operation or foods are exposed.
  - Do not apply this spray to surfaces or utensils that may come in contact with food.
- For control of insects, indoors and outdoors, in the following food and non-food areas such as:** Homes, Schools, Warehouses, Office Buildings, Apartment Buildings, Theaters, Hotels, Industrial Buildings, Motels, Kennels, Livestock Housing, Food Processing Plants, Food Service Establishments, Restaurants, Supermarkets and Grocery Stores, Transportation Equipment, Truck Trailers, Railroad Cars, and Food Manufacturing and Warehousing Establishments. Also for use on Back Yards, Lawns, Trees, Ornamental Landscaping, Recreational Areas, Parks and Athletic Fields.
- Insects Controlled:**
- |                             |             |            |                   |
|-----------------------------|-------------|------------|-------------------|
| Ants (excluding Fire Ants)  | Cockroaches | Lice       | Scorpions         |
| Aphids                      | Crickets    | Mealybugs  | Silverfish        |
| Asian Lady Beetles          | Dog Ticks   | Midges     | Spiders           |
| Bed Bugs                    | Earwigs     | Millipedes | Springtails       |
| Beetles                     | Firebrats   | Mites      | Swarming Termites |
| Boxelder Bugs               | Fleas       | Mosquitoes | Ticks             |
| Brown Marmorated Stink Bugs | Gnats       | Pillbugs   | Wasps             |
| Carpet Beetles              | Hornets     | Psocids    | Waterbugs         |
| Centipedes                  | Kudzu Bugs  | Scales     |                   |
- Flying Insects:**
- |               |             |              |                    |
|---------------|-------------|--------------|--------------------|
| Blow Flies    | Crane Flies | Fungus Gnats | Mosquitoes         |
| Chiggers      | Deer Flies  | Gnats        | Small Flying Moths |
| Clothes Moths | Face Flies  | Horn Flies   | Whiteflies         |
| Cluster Flies | Fruit Flies | House Flies  |                    |
- Biting and Stinging Pests:**
- |              |            |            |               |
|--------------|------------|------------|---------------|
| Bed Bugs     | Deer Ticks | Hornets    | Spiders       |
| Bees         | Dog Ticks  | Lice       | Ticks         |
| Biting Flies | Fleas      | Mosquitoes | Wasps         |
| Chiggers     | Gnats      | Scorpions  | Yellowjackets |
| Deer Flies   |            |            |               |
- Stored Product Pests:**
- |                        |                     |                           |                             |
|------------------------|---------------------|---------------------------|-----------------------------|
| Angoumois Grain Moths  | Flat Grain Beetles  | Meal Moth Larvae          | Spider Beetles              |
| Ants                   | Fruit Flies         | Mediterranean Flour Moths | Spider Mites                |
| Caddisflies            | Grain Mites         | Merchant Grain Beetles    | Square-necked Grain Beetles |
| Cigarette Beetles      | Grain Moths         | Red Flour Beetles         | Tobacco Moths               |
| Confused Flour Beetles | Granary Weevils     | Rice Weevils              | Yellow Mealworms            |
| Dark Mealworms         | Indian Meal Moths   | Rusty Grain Beetles       |                             |
| Driedfruit Beetles     | Lesser Grain Borers | Sawtoothed Grain Beetles  |                             |
| Drugstore Beetles      | Maize Weevils       | Skippers                  |                             |

**Livestock Pre**  
 Bed Bugs  
 Carrion Beetles  
 Chiggers  
 Darkling Beetle  
 (lesser mealworm)  
**Wood-Destro**  
 Carpenter Ants  
 Carpenter Bees  
**Plant Pests:**  
 Alfalfa Caterpillars  
 American Plum  
 Annual Bluegrass  
 Aphids  
 Apple Maggots  
 Armyworms  
 Artichoke Plum  
 Spiders  
 Bagworms  
 Balsam Woolly  
 Beet Armyworm  
 Billbugs  
 Blueberry Span  
 Brown Marmor  
 Cherry Fruit Fly  
 Cherry Fruitworm  
 Chinch Bugs  
 Codling Moths  
 Cowpea Curculionid  
 Cranberry Fruit  
 ONSLAUGHT®  
 ONSLAUGHT N  
 wall voids, or a  
 may be sprayed  
**APPLICATION**  
 Application is p  
 Plants, Grocery  
 is present. Foo  
 and dairies. Se  
 garbage rooms  
**General Surf:**  
 operation or for  
 bakeries, and o  
 with potable wa  
**Spot or Crack**  
 treated. Do not a

with soap and after use with	<b>Livestock Premise Pests:</b>	Deer Flies	Hide Beetles	Litter Beetles
	Bed Bugs	Face Flies	Horn Flies	Mites
s, Apartment upermarkets Back Yards,	<b>Wood-Destroying Pests:</b>	Deathwatch Beetles	Old House Borers	Roundheaded House Borers
	Carpenter Ants	Furniture Beetles	Powder Post Beetles	Swarming Termites
	<b>Plant Pests:</b>			
	Alfalfa Caterpillars	Cranberry Weevils	Leafhoppers	Pear Slugs
	American Plum Borers	Cucumber Beetles	Leafminers	Pecan Leaf Phylloxera
	Annual Bluegrass Weevils	Cutworms	Leafrollers	Pecan Nut Casebearers
	Aphids	Diamondback Moths	Leaf-tiers	Pecan Spittlebugs
	Apple Maggots	Elm Leaf Beetles	Lesser Appleworms	Pecan Stem Phylloxera
	Armyworms	European Pine Sawflies	Lesser Peachtree Borers	Pecan Weevils
	Artichoke Plume Moth	Fall Webworms	Loopers	Pepper Weevils
	Bagworms	Filbertworms	Lygus Bugs	Periodical Cicadas
	Balsam Woolly Adelgids	Flea Beetles	Mexican Bean Beetles	Pickleworms
	Beet Armyworms	Grasshoppers	Mites	Pilgrims
	Billbugs	Green Cloverworms	Mole Crickets	Pine Chafers
	Blueberry Spanworms	Green Fruitworms	Naval Orangeworms	Pine Coreid Bugs
	Brown Marmorated Stink Bugs	Gypsy Moth Caterpillars	Northern Pine Weevils	Plant Bugs
	Cherry Fruit Flies	Hickory Shuckworms	Oriental Fruit Moths	Plum Curculios
	Cherry Fruitworms	Imported Cabbageworms	Painted Lady Caterpillars	Red Pine Sawflies
	Chinch Bugs	Japanese Beetles	Pea Weevils	Redheaded Pine Sawflies
	Codling Moths	Kudzu Bugs	Peach Twig Borers	Red-striped Fireworms
	Cowpea Curculios	Lace Bugs	Peachtree Borers	Rindworms
	Cranberry Fruitworms	Leaf Feeding Caterpillars	Pear Psyllids	Salt Marsh Caterpillars

Mix ONSLAUGHT Insecticide before ONSLAUGHT Mi dispersion. If spr mix containing 0 a maintenance cc excessively; dam

**Formula for de** that is in the spr

(6.4) x (num Microencapsu (Gallons of in

**Tank Mixing:** ONSLAUGHT Mi currently register Do not tank mix

**INDOOR USE:**

**Application Ra**

<b>Pests</b>
--------------

**Listed Craw Insects**

**Listed Flyi Insects**

**Listed Crawlin**

Apply as a cours closets, walls, ut areas behind pip than one (1) app

**For Ant Contro**

Apply to ant trails sources of food t

**Listed Flying I**

Treat insect resti beginning of fly s

**Listed Stingin**

Apply spray to nt doors and windo

**For Control of**

Thoroughly apply and localized are for use on anima

## PRODUCT INFORMATION

ONSLAUGHT® Microencapsulated Insecticide is a microencapsulated suspension concentrate containing 0.54 pound active ingredient per gallon.

ONSLAUGHT Microencapsulated Insecticide is designed to provide long residual activity against pests listed on this label when applied as surface or spot treatments, injected into wall voids, or as a crack and crevice spray in and around residential and commercial structures, as well as on turf and landscaping. ONSLAUGHT Microencapsulated Insecticide may be sprayed on any surface that will not be stained by water.

### APPLICATION IN FOOD PROCESSING/HANDLING ESTABLISHMENTS:

Application is permitted within food and non-food areas of Food Service and Handling Establishments (places other than private residences) including: Restaurants, Meat Processing Plants, Grocery Stores, Bakeries, Food Manufacturing and Processing Establishments, and Food Warehousing Establishments. Do not treat establishments where livestock feed is present. Food areas include areas for receiving, storing, packing, canning, bottling, wrapping, boxing, preparing, edible waste storage, and enclosed processing systems, mills, and dairies. Serving areas are places where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held. Non-food areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, garages, mop closets, and storage after canning or bottling.

**General Surface Applications:** Do not apply ONSLAUGHT Microencapsulated Insecticide as a general surface application in food areas of the facility when the facility is in operation or food is exposed. Do not apply directly to food. Cover or remove all food and food processing equipment prior to application. After spraying in meat packing plants, bakeries, and other food processing plants, wash with an effective cleaning compound, and then rinse all equipment, benches, shelving, etc. where exposed food will be handled with potable water. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

**Spot or Crack and Crevice Applications:** Spot or crack and crevice applications may be made while facility is in operation, provided exposed food is covered or removed from the area being treated. Do not apply directly to food or food-handling surfaces. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

ettes

### MIXING INSTRUCTIONS

Mix ONSLAUGHT Microencapsulated Insecticide with water and apply with hand pressurized or power operated sprayers. Shake container of ONSLAUGHT Microencapsulated Insecticide before diluting. Clean spray equipment before use. For dilution, add approximately half the required water to spray tank and then add the appropriate amount of ONSLAUGHT Microencapsulated Insecticide. Agitate, and slowly add the remaining water. Agitate spray thoroughly before using and also occasionally during use to ensure dispersion. If spray filter screens are used, they should be 50 mesh or larger. Use 0.5 fl. oz. of ONSLAUGHT Microencapsulated Insecticide in 1 gallon of water to make a spray mix containing 0.025% active ingredient. Use 1.0 fl. oz. per gallon of water to make a spray containing 0.05% active ingredient. Use 0.025% solution for light infestations or as a maintenance control rate. Use 0.05% for heavy infestations or as an initial clean-out rate. Apply two-second bursts of spray per square foot of area being treated. Do not apply excessively; dampen surfaces but not to the point of saturation or runoff. Only dilute ONSLAUGHT Microencapsulated Insecticide with water.

**Formula for determining the active ingredient content of the finished spray mixture:** The following formula may be used to determine the percent active ingredient that is in the sprayer tank after mixing ONSLAUGHT Microencapsulated Insecticide:

$$\frac{(6.4) \times (\text{number of fl. oz. of ONSLAUGHT Microencapsulated Insecticide added to tank})}{(\text{Gallons of finished spray mix}) \times (128)} = \text{Percent Active Ingredient of Spray Mix}$$

**Tank Mixing:**

ONSLAUGHT Microencapsulated Insecticide may be tank mixed with an insect growth regulator such as NyGuard® IGR Concentrate or pyrethrum-containing products or any other currently registered pesticides unless expressly prohibited by the product label. The resulting tank mix may be applied in areas where these products are allowed to be sprayed. Do not tank mix ONSLAUGHT Microencapsulated Insecticide with products containing dichlorvos (DDVP).

**INDOOR USE:**

**Application Rates for Indoor Structural Pests:**

Pests	Light Infestation		Heavy Infestation		Application Rate of Mixed Solution	Application Methods
	% Active Ingredient	Dilution Rate in Water	% Active Ingredient	Dilution Rate in Water		
Listed Crawling Insects	0.025%	0.5 fl. oz./gallon	0.050%	1.0 fl. oz./ gallon	1 gallon/1,000 sq. ft.	Apply as a broadcast surface spray, crack and crevice treatment, spot treatment, or injection into wall voids.
Listed Flying Insects						

**Listed Crawling Insect Pests Indoors:**

Apply as a course, wet spray to surfaces where these pests are normally found. Treat floors, baseboards, around doors and windows, in attics, crawl spaces, eaves, corners, closets, walls, utility pipes, storage areas, and all cracks and crevices. Treat underneath sinks, dishwashers, refrigerators, stoves, the underside of shelves, drawers, cabinets, areas behind pipes, and in all places where these insects shelter. Contact as many insects as possible with direct spray. Repeat applications as needed, but do not exceed more than one (1) application every seven (7) days.

**For Ant Control Indoors:**

Apply to ant trails, around garbage receptacles, and near food sources. Also apply around doors and windows and wherever these pests may find entrance to the structure. Remove sources of food through basic hygiene practices whenever possible.

**Listed Flying Insect Pests Indoors:**

Treat insect resting areas such as walls and ceilings, screens, around windows, doors, and light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season, and repeat treatments during periods of heavy infestation.

**Listed Stinging Insect Pests Indoors:**

Apply spray to nests late in the evening when stinging insects are at rest. Thoroughly spray nest, nest entrance, and surrounding areas where insects land or walk. Treat around doors and windows, in attics, crawl spaces, and possible harborage sites or points of entry.

**For Control of Brown Dog Ticks:**

Thoroughly apply as a spot treatment to infested areas such as pet beds and resting areas, nearby cracks and crevices, along and behind baseboards, window and door frames, and localized areas of floor and floor coverings where these pests may be present. DO NOT SPRAY PETS WITH THIS PRODUCT. Treat dogs and cats with a product registered for use on animals.

ers

gs

is  
Aoths  
illars

orms

nected into  
! Insecticide

Processing  
restock feed  
terns, mills,  
reas include  
otting.

facility is in  
king plants,  
be handled

re area being

**For Control of Fleas:**

Thoroughly apply as a fine particle broadcast spray to infested rugs, carpets, and pet resting areas. Prior to treatment, cover aquariums and fish bowls. Remove animals from the area being treated. Do not permit humans or pets to contact treated surfaces until spray has dried. Remove and replace pet bedding with clean, fresh bedding after treatment. DO NOT SPRAY PETS WITH THIS PRODUCT. Treat dogs and cats with a product registered for use on animals.

**To Control Bed Bugs:**

Thoroughly clean and sanitize mattresses and box springs. Treat mattresses and box springs with an approved pesticide, such as a pyrethrin aerosol. Apply ONSLAUGHT Microencapsulated Insecticide as a spot treatment to potential harborage sites and migration paths, and cracks and crevices, around baseboards, floorboards, headboards, and walls.

**For Control of Stored Product Pests:**

Spray thoroughly around and into floor drains, non-food conveyors, benches, pipes, pallets, moist areas, storage racks, pieces of equipment, and other areas where stored product pests may be found. Tank mix or sequential use of an insect growth regulator, such as NyGuard IGR Concentrate, is recommended to break the insect reproduction cycle. Do not apply this spray to surfaces or utensils that may come in contact with food, since excessive residues in food may result.

**To Treat Voids in Equipment and Structures:**

To kill insects harboring in wall voids and other inaccessible spaces in equipment and structures, use injection equipment designed for deep void applications. Follow the injection equipment manufacturer's recommendations for proper set up and air pressures. Place the applicator tip at or into the void space to be treated. For inaccessible voids, it may be necessary to drill an access hole(s). Inject product into the void space in short bursts, allowing air pressure to push insecticide deep into the space. Do not apply to the point of runoff or drip. Ventilate area thoroughly before re-entry. Do not reapply more than every fourteen (14) days.

**To Kill the Accessible Stages of Listed Granary Insects:** Use this concentrate to treat grain storage facilities, and other listed areas, for stored product pest control. For control of exposed adult and immature stages of stored product pests, apply to cracks, crevices, and other surfaces where the pests have been seen or have harborage. Treat areas where products are stored before filling with the product. Apply at the rates listed above, using one gallon of spray mix per 1,000 square feet of surface area to be treated. Cleaning all areas prior to use will increase levels of control. Remove and destroy foodstuffs infested with pests.

Repeat application as needed, but do not exceed more than one application every fourteen (14) days.

**USE IN AND AROUND LIVESTOCK HOUSING:**

For use in and around unoccupied areas of livestock facilities, such as, but not limited to: Barns, Cow and Calf Pens and Hutches, Dairy Barns and Milk Rooms, Hog Barns, Horse Barns, Sheep Barns, Poultry Houses, and Rabbit Hutches. Do not contaminate milk, food, or drinking water. Remove animals from area being treated. Cover feeders and waterers. Do not apply or allow insecticide to drift onto animals. Do not allow animals to enter treated areas until spray solution has dried.

Pests	Light Infestation		Heavy Infestation		
	% Active Ingredient	Dilution Rate in Water	% Active Ingredient	Dilution Rate in Water	Application Methods
Listed Crawling Insects *Including Litter Beetles	0.025%	0.5 fl. oz. /gallon or sufficient water to cover 1,000 sq. ft.	0.050%	1.0 fl. oz. /gallon or sufficient water to cover 1,000 sq. ft.	Apply as a general surface spray, crack and crevice treatment, spot treatment, or injection into wall voids.
Listed Flying Insects					

**Listed Crawling Insect Pests in Livestock and Poultry Premises:**

Apply as a general surface, spot treatment, and/or crack and crevice treatment. Apply to floors and vertical and overhead surfaces where crawling insects are or may be present. Treat stanchions, pipes, windows, doors, posts, cage framing, gates, under (but not in) feeders, and other areas where insects hide or congregate. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1-3 feet up the exterior foundation wall and 3-6 feet out from the foundation.

**Litter Beetle Control in Livestock and Poultry Premises:**

To control litter beetles (darkling, hide, and carrion beetles), apply ONSLAUGHT Microencapsulated Insecticide to walls and floors at cleanout and before reintroduction of animals. Treat areas where beetles frequently occur, such as walls, supports, cages, cage framing, stalls, and around feeders. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1-3 feet up the exterior foundation wall and 3-6 feet out from the foundation.

**Listed Flying Insect Pests in Livestock and Poultry Premises:**

For residual control of listed flying insects, treat insect resting areas such walls, ceilings, screens, around windows, doors, light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season and repeat treatments during periods of heavy infestation.

**OUTDOOR USE:**

**Use Restriction:**

- Do not spray in
- Do not water the
- Do not make ap
- Do not apply dir
- Not for use on p
- All outdoor app elsewhere on th
  1. Applications
  2. Applications
  3. Applications
  4. Application to
  5. Applications
  6. Applications on this label, control occas

**Application Rat**

**Treating Exteri Foundations ar**

**Treating Nests Harborage Are Stinging Insect**

**To Control Swz Termites and V Destroying Pes**

**Treating Ornan and Landscape**

**Mosquito Bree**

**Treating Lawns**

**Treating exteri**

To control infestat carports, garages, To help prevent in exterior walls, eave Apply with sufficie per 50 gallons of v

s from the  
tment. DO

SLAUGHT  
ards, and

ad product  
le. Do not

e injection  
, it may be  
ne point of

st control.  
harborage.  
rtface area

rms, Horse  
j waterers.

ack and  
injection

sent. Treat  
ts, make a

of animals.  
r treatment

tract flying

**OUTDOOR USE:**

**Use Restrictions:**

- Do not spray in or near fish ponds or other bodies of water.
- Do not water the treated area to the point of runoff.
- Do not make applications during the rain.
- Do not apply directly to sewers or drains, or any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur, except as directed by this label.
- Not for use on plants being grown for sale or other commercial use. Not for use in nurseries, sod farms or golf courses.
- All outdoor applications, if permitted elsewhere on this label must be limited to spot or crack and crevice treatments only, except for the following permitted uses, if allowed elsewhere on this label:
  1. Applications to soil or vegetation, as listed on this label;
  2. Applications to lawns, turf, and other vegetation, as listed on this label;
  3. Applications to the side of a building, up to a maximum height of 3 feet above grade;
  4. Application to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
  5. Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width;
  6. Applications made through the use of a coarse, low pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch or other vegetation, as listed on this label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters in order to control occasional invaders or aggregating pests as listed on this label.

**Application Rates for Outdoor Pests:**

	Concentration of Active Ingredient	Dilution Rate in Water	Application Method(s)
<b>Treating Exterior Walls, Foundations and Structures</b>	0.025 to 0.050%	0.5 to 1.0 fl. oz./gallon or 0.5 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a coarse, wet spray. *General Surface Spray *Crack and Crevice Treatment *Spot Treatment
<b>Treating Nests and Harborage Areas of Listed Stinging Insects</b>	0.050%	1.0 fl. oz./gallon or 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	
<b>To Control Swarming Termites and Wood Destroying Pests</b>	0.050%	1.0 fl. oz./gallon or 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	
<b>Treating Ornamental Trees and Landscapes</b>	0.005 to 0.050%	0.1 to 1.0 fl. oz./gallon or 0.1 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a spray. *Broadcast Spray Treatment *Foliar Spray Treatment
<b>Mosquito Breeding Sites</b>	0.025%	0.5 fl. oz./gallon or 0.5 fl. oz. in sufficient water to cover 1,000 sq. ft.	
<b>Treating Lawns and Turf</b>	0.005 to 0.050%	0.1 to 1.0 fl. oz./gallon or 0.1 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft. or 4.3 to 43 fl. oz. per acre	Apply as a coarse, wet spray. *Broadcast Spray Treatment

**Treating exterior walls, foundations and structures:**

To control infestations of listed pests, treat exterior surfaces of buildings, walls, window frames, around garbage cans, eaves, cracks and crevices, porches, decks, gazebos, patios, carports, garages, fence lines, and other areas where pests are active or may be hiding.

To help prevent insect infestation of buildings: Treat a 2 to 6 foot band of soil or other substrate adjacent to buildings. Treat building foundation to a height of 2 to 3 feet. Treat exterior walls, eaves, cracks and crevices, and other areas where listed pests are active, and may find entrance into building.

Apply with sufficient water to adequately cover the area being treated, but do not allow dripping or runoff to occur. Alternate mixing directions are to use 2 to 4 fl. oz. of concentrate per 50 gallons of water.

**Listed Stinging I**

For listed stinging ins and surrounding areas pests are active, appli For Yellowjacket con

**Insect Control in**

For residential and c appropriate volumes to surrounding soil, but no more than or Spray and observe f

**Insect Control on**

For best results, law 1,000 sq. ft. to get u with at least 0.25 inc When hand spraying trees. Repeat treatm

**Rate Range:**

Use lower rate rang at the time of applic

**To Kill Swarming**

Apply spray mix as usually occurs in the colonies. This treatr (winged) and worke

**To Kill Listed Wc**

Apply spray mix as window and door fra wooden structural vi

**Mosquito Breedi**

Mosquito populatio into tall grass, shru Not for wide area m

Do not contaminate

**PESTICIDE STOF**

pet food, feed, seec

**PESTICIDE DISP**

to a waste disposa

**CONTAINER HAN**

Non-refillable cont

Triple rinse as foll

1/4 full with water

after the flow begir

Do not burn unles

MGK®, ONSLAUGHT

his label.

allowed

as listed  
order to

as listed  
order to

s, patios,

et. Treat

ncentrate

**Listed Stinging Insect Control Outdoors:**

For listed stinging insects, apply spray solution to nests and harborage areas late in the evening when stinging insects are commonly at rest. Thoroughly spray nest, nest entrance, harborage sites, and surrounding areas where insects land or walk. For nests inside walls, inject sufficient spray to treat nest. Do not use in areas where an electrical shock hazard exists. For applications made when pests are active, applicator should wear protective equipment as required. Repeat application when there are signs of renewed insect activity.

For Yellowjacket control, ONSLAUGHT Microencapsulated Insecticide can be mixed with baits in traps. Follow trap instructions for preparation of bait.

**Insect Control in Ornamental Trees and Landscapes:**

For residential and commercial trees, shrubs, ground covers, and bedding and foliage plants that will not be harvested for food, apply ONSLAUGHT Microencapsulated Insecticide in appropriate volumes of water to obtain thorough coverage. Apply as a full-coverage foliar spray, applying to the point of drip but not runoff. Treat active soil pests with an application to surrounding soil. Repeat treatments as necessary to achieve control, using higher application rates as pest pressure and foliage area increases. Repeat applications as necessary, but no more than once every seven (7) days. Certain plants may be sensitive to the final spray solution. A small-scale test is recommended to verify safety to ornamental plants. Spray and observe for one (1) week prior to application of an entire planting.

**Insect Control on Lawns, Turf Grass, and Turf:**

For best results, lawn or turf grass should be mowed 1-2 days before spraying. Treat with spraying equipment or a hand sprayer. Use application volumes of up to 10 gallons per 1,000 sq. ft. to get uniform coverage when treating dense grass foliage. For low volume applications using less than 2 gallons of spray solution per 1,000 sq. ft., immediate irrigation with at least 0.25 inch of water is recommended to improve effectiveness on sub-surface pests.

When hand spraying, spray using a slow, even sweeping motion, making sure to cover the entire lawn or turf grass area where pests are observed. Spray under ornamentals and trees. Repeat treatments may be necessary at 7 to 14 day intervals.

**Rate Range:**

Use lower rate range for pests that are commonly exposed and will be contacted by spray solution at the time of application. For pests that will not be contacted by spray solution at the time of application, use the upper rate range. Use 1.0 fl. oz. per 1,000 sq. ft. when treating Mole Crickets and Chinch Bugs.

**To Kill Swarming Termites: OUTDOORS ONLY.**

Apply spray mix as a coarse, wetting spray when swarming termites are seen emerging from woodpiles, wooden fence posts, wooden structures, or from the ground. Swarming usually occurs in the spring or at other times when a termite colony becomes overcrowded and new reproductive termites with wings emerge and fly away to mate and establish new colonies. This treatment will control the sprayed termites and will not protect the structure from which the swarm is coming. Use only as a contact spray to kill emerging reproductive (winged) and worker termites emerging from infested wood. This treatment is not a substitute for a comprehensive termite control program.

**To Kill Listed Wood Destroying Pests: OUTDOORS ONLY.**

Apply spray mix as a coarse, wetting spray to exposed pests and to the damaged areas of wood, spraying into galleries or tunnels in the exposed wood. Also spray around doors, window and door frames, and other areas where these pests may hide or enter the house. Spray into cracks and crevices and, if necessary, drill small holes and spray into inaccessible wooden structural voids where these pests are suspected.

**Mosquito Breeding sites:**

Mosquito populations may be reduced by application of ONSLAUGHT Microencapsulated Insecticide to sites where mosquitoes rest, harbor, and breed. Apply spray solution into tall grass, shrubbery, and around back yards and lawns where these pests may hover or rest. Apply while air is still.

Not for wide area mosquito abatement using aerial or truck mounted cold aerosol ULV sprayers and thermal fogging devices.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local government or by industry).

**CONTAINER HANDLING**

Non-refillable container. Do not reuse or refill this container.

Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into the application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

MGK®, ONSLAUGHT® and NyGuard® are registered trademarks of McLaughlin Gormley King Company



**Mic**

**ACTIVE ING**  
(S)-cyano(3-  
\*OTHER INGRI  
\*Contains petr

**See inside b**

Manufactured fr  
EPA Reg. No.  
F28371-107  
Rev. 0616-101







# Microencapsulated Insecticide

**ACTIVE INGREDIENT:**

(S)-cyano(3-phenoxyphenyl)methyl-(S)-4-chloro-alpha-(1-methylethyl)benzeneacetate.....	6.40%
*OTHER INGREDIENTS.....	93.60%
*Contains petroleum distillates	100.00%

## KEEP OUT OF REACH OF CHILDREN CAUTION

**See inside booklet for First Aid, Precautionary Statements, Directions for Use and Storage and Disposal**

Manufactured for: **MGK**, 8810 Tenth Avenue North, Minneapolis, MN 55427

EPA Reg. No. 1021-1815  
F28371-107

Rev. 0616-1017

EPA Est. No. 1021-MN-2<sup>a</sup>, 70815-GA-1<sup>c</sup>, 58996-MO-1<sup>d</sup>,  
39578-TX-1<sup>e</sup>, 228-IL-002<sup>g</sup>, 10508-OH-001<sup>k</sup>, 67572-GA-1<sup>l</sup>,

63416-WI-001<sup>m</sup>, 5905-GA-01<sup>o</sup>

Superscript is the first character of the lot number.

**NET CONTENTS: 1 GALLON**

ge sites,  
Je when

icide in  
lication  
essary,  
plants.

ons per  
tigation

als and

olution

arming  
sh new  
ductive

doors,  
essible

olution

iod,

luct

iner  
nds  
ion.

ACCEPTED  
VIA NOTIFICATION  
LABEL NOT REVIEWED

March 9, 2018

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration



**Microencapsulated  
Insecticide**

**ACTIVE INGREDIENT:**  
(S)-cyano(3-phenoxyphenyl)  
methyl-(S)-4-chloro-alpha-  
(1-methylethyl)benzeneacetate . . . 6.40%  
\*OTHER INGREDIENTS . . . . . 93.60%  
100.00%

\*Contains petroleum distillates.

**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

**FIRST AID**  
**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

Manufactured for:  
**MGK**, 8810 Tenth Avenue North, Minneapolis, MN 55427  
EPA Reg. No. 1021-1815 EPA Est. No. 1021-MN-2<sup>a</sup>, 70815-GA-1<sup>c</sup>,  
F28371-005 58996-MO-1<sup>b</sup>, 39578-TX-1<sup>f</sup>, 228-IL-002<sup>e</sup>,  
Rev. 0616-1017 10508-OH-001<sup>k</sup>, 67572-GA-1<sup>l</sup>,  
63416-WI-001<sup>m</sup>, 5905-GA-01<sup>o</sup>  
Superscript is the first character  
of the lot number.

**NET CONTENTS: 1 PINT**

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**NOTE TO PHYSICIAN:**  
Contains petroleum distillate - vomiting may cause aspiration pneumonia.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

**See inside booklet for Additional Precautionary Statements, Directions for Use and Storage and Disposal**

PULL HERE TO OPEN / PRESS TO RESEAL

PF 20037-14

FIRST AID	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not give <b>any</b> liquid to the person.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or a doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>NOTE TO PHYSICIAN:</b> Contains petroleum distillate – vomiting may cause aspiration pneumonia.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear protective eyewear (goggles, face shield or safety glasses with side shields). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and other aquatic invertebrates. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage system. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirement of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

This product is highly toxic to bees and other pollinating insects exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees or other pollinating insects are foraging in the treatment area.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS:**

- Remove pets and birds and cover fish aquariums before spraying.
- Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm.
- Do not apply to classrooms while in use. Do not apply in institutions (including libraries, schools, sports facilities, etc.) in the immediate area where occupants are present.
- DO NOT use this product as a fogger or apply it as a space spray.
- Do not apply in electrical conduits, junction and switch boxes, motor housings, or other electrical equipment due to shock hazard from water-based spray.
- Use only in areas described.
- Remove pets and cover fish aquariums and pet's food and water dishes before spraying.
- Keep all people (especially children) and pets out of areas being treated, and restrict access to these areas until all surfaces are dry.
- DO NOT leave product where children or pets can come into contact with it.
- DO NOT allow spray to contact food or food-containing surfaces, feed, or water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated with this product. Food utensils such as teaspoons and measuring cups must not be used for food purposes after use with pesticides.
- Do not apply when food processing facility is in operation or foods are exposed.
- Do not apply this spray to surfaces or utensils that may come in contact with food.

**For control of insects, indoors and outdoors, in the following food and non-food areas such as:** Homes, Schools, Warehouses, Office Buildings, Apartment Buildings, Theaters, Hotels, Industrial Buildings, Motels, Kennels, Livestock Housing, Food Processing Plants, Food Service Establishments, Restaurants, Supermarkets and Grocery Stores, Transportation Equipment, Truck Trailers, Railroad Cars, and Food Manufacturing and Warehousing Establishments. Also for use on Back Yards, Lawns, Trees, Ornamental Landscaping, Recreational Areas, Parks and Athletic Fields.

**Insects Controlled:**

Ants (excluding Fire Ants)	Centipedes	Kudzu Bugs	Scorpions
Aphids	Cockroaches	Lice	Silverfish
Asian Lady Beetles	Crickets	Mealybugs	Spiders
Bed Bugs	Dog Ticks	Midges	Springtails
Beetles	Earwigs	Millipedes	Swarming Termites
Boxelder Bugs	Firebrats	Mites	Ticks
Brown Marmorated Stink Bugs	Fleas	Mosquitoes	Wasps
Carpet Beetles	Gnats	Pillbugs	Waterbugs
	Hornets	Psocids	

**Flyin**  
Blow  
Chigg  
Cloth  
Clust

**Bitin**  
Bed E  
Bees  
Biting  
Chigg  
Deer

**Storr**  
Ango  
Ants  
Cadel  
Ciger  
Conit  
Dark  
Dried  
Drugs

**Live:**  
Bed E  
Carrir  
Chigg  
Darkl  
(les

**Woo**  
Carpe  
Carpe

	<p><b>Flying Insects:</b> Blow Flies Chiggers Clothes Moths Cluster Flies</p> <p><b>Biting and Stinging Pests:</b> Bed Bugs Bees Biting Flies Chiggers Deer Flies</p> <p><b>Stored Product Pests:</b> Angoumois Grain Moths Ants Cadelles Cigarette Beetles Confused Flour Beetles Dark Mealworms Driedfruit Beetles Drugstore Beetles</p> <p><b>Livestock Premise Pests:</b> Bed Bugs Carrion Beetles Chiggers Darkling Beetles (lesser mealworms)</p> <p><b>Wood-Destroying Pests:</b> Carpenter Ants Carpenter Bees</p>	<p>Crane Flies Deer Flies Face Flies Fruit Flies</p> <p>Deer Ticks Dog Ticks Fleas Gnats Hornets</p> <p>Flat Grain Beetles Fruit Flies Grain Mites Grain Moths Granary Weevils Indian Meal Moths Lesser Grain Borers Maize Weevils</p> <p>Deer Flies Face Flies Fleas Flies Hide Beetles</p> <p>Deathwatch Beetles Furniture Beetles</p>	<p>Fungus Gnats Gnats Horn Flies House Flies</p> <p>Lice Mosquitoes Scorpions Spiders Ticks</p> <p>Meal Moth Larvae Mediterranean Flour Moths Merchant Grain Beetles Red Flour Beetles Rice Weevils Rusty Grain Beetles Sawtoothed Grain Beetles Skippers</p> <p>Horn Flies Horse Flies Lice Litter Beetles Mites</p> <p>Old House Borers Powder Post Beetles</p>	<p>Mosquitoes Small Flying Moths Whiteflies</p> <p>Wasps Yellowjackets</p> <p>Spider Beetles Spider Mites Squarenecked Grain Beetles Tobacco Moths Yellow Mealworms</p> <p>Mosquitoes Stable Flies</p> <p>Roundheaded House Borers Swarming Termites</p>	<p><b>Plant Pests:</b> Ailanthus Caterpillars American Plum Borers Annual Bluegrass Weevils Aphids Apple Maggots Armyworms Artichoke Plume Moth Bagworms Balsam Woolly Adelgids Beet Armyworms Billbugs Blueberry Spanworms Brown Marmorated Stink Bugs Cherry Fruit Flies Cherry Fruitworms Chinch Bugs Coddling Moths Cowpea Curculios Cranberry Fruitworms Cranberry Weevils Cucumber Beetles Cutworms Diamondback Moths</p> <p>Elm Leaf Beetles European Pine Sawflies Fall Webworms Filbertworms Flea Beetles Grasshoppers Green Cloverworms Green Fruitworms Gypsy Moth Caterpillars Hickory Shuckworms Imported Cabbageworms Japanese Beetles Kudzu Bugs Lace Bugs Leaf Feeding Caterpillars Leathoppers Leafminers Leafrollers Leafhoppers Lesser Appleworms Lesser Peachtree Borers Loopers Lygus Bugs</p> <p>Mexican Bean Beetles Mites Mole Crickets Naval Orangeworms Northern Pine Weevils Oriental Fruit Moths Painted Lady Caterpillars Pea Weevils Peach Twig Borers Peachtree Borers Pear Psyllids Pear Slugs Pecan Leaf Phylloxera Pecan Nut Casebearers Pecan Spittlebugs Pecan Stem Phylloxera Pecan Weevils Pepper Weevils Periodical Cicadas Pickleworms Pillbugs Pine Chafers Pine Coreid Bugs</p>	<p>Plant Bugs Plum Curculios Red Pine Sawflies Redheaded Pine Sawflies Red-striped Fireworms Rindworms Salt Marsh Caterpillars Sap Beetles Scales Sod Webworms Sowbugs Spider Slugs Spittlebugs Stink Bugs Tarnished Plant Bugs Tent Caterpillars Thrips Tobacco Hornworms Tufted Apple Bud Moths Velvet Bean Caterpillars Walnut Husk Flies Western Bean Cutworms and Whiteflies</p>
<p>facilities, etc.) in the</p> <p>it due to shock hazard</p> <p>reas until all surfaces</p> <p>ash dishes and food ch as teaspoons and</p> <p>as: Homes, Schools, , Livestock Housing, portation Equipment, ; Yards, Lawns, Trees,</p> <p>Termites</p>						

## PRODUCT INFORMATION

ONSLAUGHT® Microencapsulated Insecticide is a microencapsulated suspension concentrate containing 0.54 pound active ingredient per gallon.

ONSLAUGHT Microencapsulated Insecticide is designed to provide long residual activity against pests listed on this label when applied as surface or spot treatments, injected into wall voids, or as a crack and crevice spray in and around residential and commercial structures, as well as on turf and landscaping. ONSLAUGHT Microencapsulated Insecticide may be sprayed on any surface that will not be stained by water.

### APPLICATION IN FOOD PROCESSING/HANDLING ESTABLISHMENTS:

Application is permitted within food and non-food areas of Food Service and Handling Establishments (places other than private residences) including: Restaurants, Meat Processing Plants, Grocery Stores, Bakeries, Food Manufacturing and Processing Establishments, and Food Warehousing Establishments. Do not treat establishments where livestock feed is present. Food areas include areas for receiving, storing, packing, canning, bottling, wrapping, boxing, preparing, edible waste storage, and enclosed processing systems, mills, and dairies. Serving areas are places where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held. Non-food areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, garages, mop closets, and storage after canning or bottling.

ant Bugs  
um Curculios  
d Pine Sawflies  
d-headed Pine Sawflies  
d-striped Fireworms  
ndworms  
lt Marsh Caterpillars  
p Beetles  
ales  
d Webworms  
wbugs  
iders  
ittlebugs  
ink Bugs  
rnished Plant Bugs  
nt Caterpillars  
rips  
bacco Hornworms  
fted Apple Bud Moths  
lvet Bean Caterpillars  
alnut Husk Flies  
estern Bean Cutworms  
and Whiteflies

und active ingredient per gallon.  
this label when applied as surface  
rcial structures, as well as on turf  
ined by water.

er than private residences) includ-  
ishments, and Food Warehousing  
iving, storing, packing, canning,  
s. Serving areas are places where  
Non-food areas include garbage  
s, mop closets, and storage after

**General Surface Applications:** Do not apply ONSLAUGHT Microencapsulated Insecticide as a general surface application in food areas of the facility when the facility is in operation or food is exposed. Do not apply directly to food. Cover or remove all food and food processing equipment prior to application. After spraying in meat packing plants, bakeries, and other food processing plants, wash with an effective cleaning compound, and then rinse all equipment, benches, shelving, etc. where exposed food will be handled with potable water. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

**Spot or Crack and Crevice Applications:** Spot or crack and crevice applications may be made while facility is in operation, provided exposed food is covered or removed from the area being treated. Do not apply directly to food or food-handling surfaces. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

### MIXING INSTRUCTIONS

Mix ONSLAUGHT Microencapsulated Insecticide with water and apply with hand pressurized or power operated sprayers. Shake container of ONSLAUGHT Microencapsulated Insecticide before diluting. Clean spray equipment before use. For dilution, add approximately half the required water to spray tank and then add the appropriate amount of ONSLAUGHT Microencapsulated Insecticide. Agitate, and slowly add the remaining water. Agitate spray thoroughly before using and also occasionally during use to ensure dispersion. If spray filter screens are used, they should be 50 mesh or larger. Use 0.5 fl. oz. of ONSLAUGHT Microencapsulated Insecticide in 1 gallon of water to make a spray mix containing 0.025% active ingredient. Use 1.0 fl. oz. per gallon of water to make a spray containing 0.05% active ingredient. Use 0.025% solution for light infestations or as a maintenance control rate. Use 0.05% for heavy infestations or as an initial clean-out rate. Apply two-second bursts of spray per square foot of area being treated. Do not apply excessively; dampen surfaces but not to the point of saturation or runoff. Only dilute ONSLAUGHT Microencapsulated Insecticide with water.

**Formula for determining the active ingredient content of the finished spray mixture:** The following formula may be used to determine the percent active ingredient that is in the sprayer tank after mixing ONSLAUGHT Microencapsulated Insecticide:

$$\frac{(6.4) \times (\text{number of fl. oz. of ONSLAUGHT Microencapsulated Insecticide added to tank})}{(\text{Gallons of finished spray mix}) \times (128)} = \text{Percent Active Ingredient of Spray Mix}$$

**Tank Mixing:**

ONSLAUGHT Microencapsulated Insecticide may be tank mixed with an insect growth regulator such as NyGuard® IGR Concentrate or pyrethrum-containing products or any other currently registered pesticides unless expressly prohibited by the product label. The resulting tank mix may be applied in areas where these products are allowed to be sprayed. Do not tank mix ONSLAUGHT Microencapsulated Insecticide with products containing dichlorvos (DDVP).

**INDOOR USE:**

**Application Rates for Indoor Structural Pests:**

Pests	Light Infestation		Heavy Infestation		Application Rate of Mixed Solution	Application Methods
	% Active Ingredient	Dilution Rate in Water	% Active Ingredient	Dilution Rate in Water		
Listed Crawling Insects	0.025%	0.5 fl. oz./gallon	0.050%	1.0 fl. oz./gallon	1 gallon/1,000 sq. ft.	Apply as a broadcast surface spray, crack and crevice treatment, spot treatment, or injection into wall voids.
Listed Flying Insects						

**Listed Crawling Insect Pests Indoors:**

Apply as a course, wet spray to surfaces where these pests are normally found. Treat floors, baseboards, around doors and windows, in attics, crawl spaces, eaves, corners, closets, walls, utility pipes, storage areas, and all cracks and crevices. Treat underneath sinks, dishwashers, refrigerators, stoves, the underside of shelves, drawers, cabinets, areas behind pipes, and in all places where these insects shelter. Contact as many insects as possible with direct spray. Repeat applications as needed, but do not exceed more than one (1) application every seven (7) days.

**For Ant Control Indoors:**

Apply to ant trails, around garbage receptacles, and near food sources. Also apply around doors and windows and wherever these pests may find entrance to the structure. Remove sources of food through basic hygiene practices whenever possible.

**Listed Flying Insect Pests Indoors:**

Treat insect resting areas such as walls and ceilings, screens, around windows, doors, and light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season, and repeat treatments during periods of heavy infestation.

**Listed Stinging Insect Pests Indoors:**

Apply spray to nests late in the evening when stinging insects are at rest. Thoroughly spray nest, nest entrance, and surrounding areas where insects land or walk. Treat around doors and windows, in attics, crawl spaces, and possible harborage sites or points of entry.

**For Control of Brown Dog Ticks:**

Thoroughly apply as a spot treatment to infested areas such as pet beds and resting areas, nearby cracks and crevices, along and behind baseboards, window and door frames, and localized areas of floor and floor coverings where these pests may be present. DO NOT SPRAY PETS WITH THIS PRODUCT. Treat dogs and cats with a product registered for use on animals.

**For Control of Fleas:**

Thoroughly apply as a fine particle broadcast spray to infested rugs, carpets, and pet resting areas. Prior to treatment, cover aquariums and fish bowls. Remove animals from the area being treated. Do not permit humans or pets to contact treated surfaces until spray has dried. Remove and replace pet bedding with clean, fresh bedding after treatment. DO NOT SPRAY PETS WITH THIS PRODUCT. Treat dogs and cats with a product registered for use on animals.

**To Control Bed Bugs:**

Thoroughly clean and sanitize mattresses and box springs. Treat mattresses and box springs with an approved pesticide, such as a pyrethrin aerosol. Apply ONSLAUGHT Microencapsulated Insecticide as a spot treatment to potential harborage sites and migration paths, and cracks and crevices, around baseboards, floorboards, headboards, and walls.

**For Control of Stored Product Pests:**

Spray thoroughly around and into floor drains, non-food conveyors, benches, pipes, pallets, moist areas, storage racks, pieces of equipment, and other areas where stored product pests may be found. Tank mix or sequential use of an insect growth regulator, such as NyGuard IGR Concentrate, is recommended to break the insect reproduction cycle. Do not apply this spray to surfaces or utensils that may come in contact with food, since excessive residues in food may result.

**To Treat Voids in Equipment and Structures:**

To kill insects harboring in wall voids and other inaccessible spaces in equipment and structures, use injection equipment designed for deep void applications. Follow the injection equipment manufacturer's recommendations for proper set up and air pressures. Place the applicator tip at or into the void space to be treated. For inaccessible voids, it may be necessary to drill an access hole(s). Inject product into the void space in short bursts, allowing air pressure to push insecticide deep into the space. Do not apply to the point of runoff or drip. Ventilate area thoroughly before re-entry. Do not reapply more than every fourteen (14) days.

**To Kill the Accessible Stages of Listed Granary Insects:** Use this concentrate to treat grain storage facilities, and other listed areas, for stored product pest control. For control of exposed adult and immature stages of stored product pests, apply to cracks, crevices, and other surfaces where the pests have been seen or have harborage. Treat areas where products are stored before filling with the product. Apply at the rates listed above, using one gallon of spray mix per 1,000 square feet of surface area to be treated. Cleaning all areas prior to use will increase levels of control. Remove and destroy foodstuffs infested with pests.

Repeat application as needed, but do not exceed more than one application every fourteen (14) days.

**USE IN AND AROUND LIVESTOCK HOUSING:**

For use in and around unoccupied areas of livestock facilities, such as, but not limited to: Barns, Cow and Calf Pens and Hutches, Dairy Barns and Milk Rooms, Hog Barns, Horse Barns, Sheep Barns, Poultry Houses, and Rabbit Hutches. Do not contaminate milk, food, or drinking water. Remove animals from area being treated. Cover feeders and waterers. Do not apply or allow insecticide to drift onto animals. Do not allow animals to enter treated areas until spray solution has dried.

Pests	Light Infestation		Heavy Infestation		Application Methods
	% Active Ingredient	Dilution Rate in Water	% Active Ingredient	Dilution Rate in Water	
Listed Crawling Insects *Including Litter Beetles	0.025%	0.5 fl. oz. /gallon or sufficient water to cover 1,000 sq. ft.	0.050%	1.0 fl. oz. /gallon or sufficient water to cover 1,000 sq. ft.	Apply as a general surface spray, crack and crevice treatment, spot treatment, or injection into wall voids.
Listed Flying Insects					

**Listed Crawling Insect Pests in Livestock and Poultry Premises:**

Apply as a general surface, spot treatment, and/or crack and crevice treatment. Apply to floors and vertical and overhead surfaces where crawling insects are or may be present. Treat stanchions, pipes, windows, doors, posts, cage framing, gates, under (but not in) feeders, and other areas where insects hide or congregate. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1-3 feet up the exterior foundation wall and 3-6 feet out from the foundation.

**Litter Beetle Control in Livestock and Poultry Premises:**

To control litter beetles (darkling, hide, and carrion beetles), apply ONSLAUGHT Microencapsulated Insecticide to walls and floors at cleanout and before reintroduction of animals. Treat areas where beetles frequently occur, such as walls, supports, cages, cage framing, stalls, and around feeders. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1-3 feet up the exterior foundation wall and 3-6 feet out from the foundation.

**Listed Flying Insect Pests in Livestock and Poultry Premises:**

For residual control of listed flying insects, treat insect resting areas such walls, ceilings, screens, around windows, doors, light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season and repeat treatments during periods of heavy infestation.

**OUTDOOR USE:**

**Use Restrictions:**

- Do not spray in or near fish ponds or other bodies of water.
- Do not water the treated area to the point of runoff.
- Do not make applications during the rain.
- Do not apply directly to sewers or drains, or any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur, except as directed by this label.
- Not for use on plants being grown for sale or other commercial use. Not for use in nurseries, sod farms or golf courses.
- All outdoor applications, if permitted elsewhere on this label must be limited to spot or crack and crevice treatments only, except for the following permitted uses, if allowed elsewhere on this label:
  1. Applications to soil or vegetation, as listed on this label;
  2. Applications to lawns, turf, and other vegetation, as listed on this label;
  3. Applications to the side of a building, up to a maximum height of 3 feet above grade;
  4. Application to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
  5. Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width;
  6. Applications made through the use of a coarse, low pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch or other vegetation, as listed on this label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters in order to control occasional invaders or aggregating pests as listed on this label.

**Applicati**

Treating Walls, Found Structural

Treating Harbora of Lister Insects

To Cont Termite Destroy

Treating Trees a

Mosquit Sites

Treating and Tur

Treating To control porches, c

To help pr height of 2

Apply with use 2 to 4

Listed Si

For listed s

ies of water.  
off.  
y area like a gutter  
odies, or aquatic  
el.  
: other commercial  
ourses.  
on this label must  
ts only, except for  
ere on this label:  
in this label;  
getation, as listed  
a maximum height  
doors, or windows  
overing, overhang,  
nts into buildings,  
one inch in width;  
arse, low pressure  
are directly above  
n, as listed on this  
drainage or other  
m drains, drainage  
control occasional  
; label.

**Application Rates for Outdoor Pests:**

	Concentration of Active Ingredient	Dilution Rate in Water	Application Method(s)
<b>Treating Exterior Walls, Foundations and Structures</b>	0.025 to 0.050%	0.5 to 1.0 fl. oz./gallon or 0.5 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a coarse, wet spray. *General Surface Spray *Crack and Crevice Treatment *Spot Treatment
<b>Treating Nests and Harborage Areas of Listed Stinging Insects</b>	0.050%	1.0 fl. oz./gallon or 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	
<b>To Control Swarming Termites and Wood Destroying Pests</b>	0.050%	1.0 fl. oz./gallon or 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	
<b>Treating Ornamental Trees and Landscapes</b>	0.005 to 0.050%	0.1 to 1.0 fl. oz./gallon or 0.1 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a spray. *Broadcast Spray Treatment *Foliar Spray Treatment
<b>Mosquito Breeding Sites</b>	0.025%	0.5 fl. oz./gallon or 0.5 fl. oz. in sufficient water to cover 1,000 sq. ft.	
<b>Treating Lawns and Turf</b>	0.005 to 0.050%	0.1 to 1.0 fl. oz./gallon or 0.1 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a coarse, wet spray. *Broadcast Spray Treatment

**Treating exterior walls, foundations and structures:**

To control infestations of listed pests, treat exterior surfaces of buildings, walls, window frames, around garbage cans, eaves, cracks and crevices, porches, decks, gazebos, patios, carports, garages, fence lines, and other areas where pests are active or may be hiding.

To help prevent insect infestation of buildings: Treat a 2 to 6 foot band of soil or other substrate adjacent to buildings. Treat building foundation to a height of 2 to 3 feet. Treat exterior walls, eaves, cracks and crevices, and other areas where listed pests are active, and may find entrance into building.

Apply with sufficient water to adequately cover the area being treated, but do not allow dripping or runoff to occur. Alternate mixing directions are to use 2 to 4 fl. oz. of concentrate per 50 gallons of water.

**Listed Stinging Insect Control Outdoors:**

For listed stinging insects, apply spray solution to nests and harborage areas late in the evening when stinging insects are commonly at rest.

Thoroughly spray nest, nest entrance, harborage sites, and surrounding areas where insects land or walk. For nests inside walls, inject sufficient spray to treat nest. Do not use in areas where an electrical shock hazard exists. For applications made when pests are active, applicator should wear protective equipment as required. Repeat application when there are signs of renewed insect activity.

For Yellowjacket control, ONSLAUGHT Microencapsulated Insecticide can be mixed with baits in traps. Follow trap instructions for preparation of bait.

**Insect Control in Ornamental Trees and Landscapes:**

For residential and commercial trees, shrubs, ground covers, and bedding and foliage plants that will not be harvested for food, apply ONSLAUGHT Microencapsulated Insecticide in appropriate volumes of water to obtain thorough coverage. Apply as a full-coverage foliar spray, applying to the point of drip but not runoff. Treat active soil pests with an application to surrounding soil. Repeat treatments as necessary to achieve control, using higher application rates as pest pressure and foliage area increases. Repeat applications as necessary, but no more than once every seven (7) days. Certain plants may be sensitive to the final spray solution. A small-scale test is recommended to verify safety to ornamental plants. Spray and observe for one (1) week prior to application of an entire planting.

**Insect Control on Lawns, Turf Grass, and Turf:**

For best results, lawn or turf grass should be mowed 1-2 days before spraying. Treat with spraying equipment or a hand sprayer. Use application volumes of up to 10 gallons per 1,000 sq. ft. to get uniform coverage when treating dense grass foliage. For low volume applications using less than 2 gallons of spray solution per 1,000 sq. ft., immediate irrigation with at least 0.25 inch of water is recommended to improve effectiveness on sub-surface pests.

When hand spraying, spray using a slow, even sweeping motion, making sure to cover the entire lawn or turf grass area where pests are observed. Spray under ornamentals and trees. Repeat treatments may be necessary at 7 to 14 day intervals.

**Rate Range:**

Use lower rate range for pests that are commonly exposed and will be contacted by spray solution at the time of application. For pests that will not be contacted by spray solution at the time of application, use the upper rate range. Use 1.0 fl. oz. per 1,000 sq. ft. when treating Mole Crickets and Chinch Bugs.

**To Kill Swarming Termites: OUTDOORS ONLY.**

Apply spray mix as a coarse, wetting spray when swarming termites are seen emerging from woodpiles, wooden fence posts, wooden structures, or from the ground. Swarming usually occurs in the spring or at other times when a termite colony becomes overcrowded and new reproductive termites with wings emerge and fly away to mate and establish new colonies. This treatment will control the sprayed termites and will not protect the structure from which the swarm is coming. Use only as a contact spray to kill emerging reproductive (winged) and worker termites emerging from infested wood. This treatment is not a substitute for a comprehensive termite control program.

**To Kill Listed Wood Destroying Pests: OUTDOORS ONLY.**

Apply spray mix as a coarse, wetting spray to exposed pests and to the damaged areas of wood, spraying into galleries or tunnels in the exposed wood. Also spray around doors, window and door frames, and other areas where these pests may hide or enter the house. Spray into cracks and crevices and, if necessary, drill small holes and spray into inaccessible wooden structural voids where these pests are suspected.

**Mosquito Breeding sites:**

Mosquito populations may be reduced by application of ONSLAUGHT Microencapsulated Insecticide to sites where mosquitoes rest, harbor, and breed. Apply spray solution into tall grass, shrubbery, and around back yards and lawns where these pests may hover or rest. Apply while air is still.

Not for wide area mosquito abatement using aerial or truck mounted cold aerosol ULV sprayers and thermal fogging devices.

**DOORS ONLY.**

Apply spray when swarming termites are present on wooden fence posts. Swarming usually occurs when a termite colony becomes crowded and new termites with wings emerge from colonies. This treatment will not protect the structure only as a contact spray to kill worker termites emerging from a substitute for a com-

**Outdoor Pests: OUTDOORS**

Apply to exposed pests and termites in galleries or tunnels under doors, window and door frames may hide or enter the structure, if necessary, drill small holes in structural voids where

Apply by application of insecticide to sites where termites are present. Apply spray solution into cracks, yards and lawns where termites are still.

Apply using aerial or truck-mounted thermal fogging

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local government or by industry).

**CONTAINER HANDLING**

Non-refillable container. Do not reuse or refill this container.

Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into the application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

MGK®, ONSLAUGHT® and NyGuard® are registered trademarks of McLaughlin Gormley King Company



**Microencapsulated Insecticide**

**ACTIVE INGREDIENT:**

(S)-cyano(3-phenoxyphenyl)methyl-(S)-4-chloro-	
alpha-(1-methylethyl)benzeneacetate.....	6.40%
*OTHER INGREDIENTS.....	93.60%
	100.00%

\*Contains petroleum distillates

**KEEP OUT OF REACH OF CHILDREN CAUTION**

See inside booklet for First Aid, Precautionary Statements, Directions for Use and Storage and Disposal

Manufactured for:

**MGK**, 8810 Tenth Avenue North, Minneapolis, MN 55427

EPA Reg. No. 1021-1815  
F28371-005  
Rev. 0616-1017

EPA Est. No. 1021-MN-2<sup>0</sup>,  
70815-GA-1<sup>0</sup>, 58996-MO-1<sup>0</sup>,  
39578-TX-1<sup>0</sup>, 228-IL-002<sup>0</sup>,  
10508-OH-001<sup>0</sup>, 67572-GA-1<sup>0</sup>,  
63416-WI-001<sup>0</sup>, 5905-GA-01<sup>0</sup>  
Superscript is the first character  
of the lot number.

**NET CONTENTS: 1 PINT**